

# Exercise Sheet 4

Discussion on 17.11.23

## Exercise 1

Prove Lemma 4.13 from the lecture:

A sequence  $(y_k)_{k=1,\dots}$  is a solution of the difference equation given by  $(\alpha_\ell)_{\ell=0,\dots,m}$  if and only if the vectors

$$Y_k = (y_k, y_{k+1}, \dots, y_{k+m-1})^\top$$

satisfy  $Y_{k+1} = AY_k$  for  $k = 0, 1, \dots$ . The companion matrix  $A \in \mathbb{R}^{m \times m}$  is defined by

$$A = \begin{pmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \\ -\alpha_0 & -\alpha_1 & \cdots & -\alpha_m \end{pmatrix}.$$

If  $A$  has linear independent eigenvectors  $v_1, v_2, \dots, v_m \in \mathbb{R}^m$  with corresponding eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_m$  and if  $\gamma_1, \gamma_2, \dots, \gamma_m \in \mathbb{R}$  are the coefficients of vector  $Y_0$  with respect to this basis,

$$Y_k = A^k Y_0 = \sum_{j=1}^m \lambda_j^k \gamma_j v_j.$$

holds.

The eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_m$  are the roots of the characteristic polynomial

$$q(\lambda) = \lambda^m + \alpha_{m-1} \lambda^{m-1} + \cdots + \lambda \alpha_1 + \alpha_0.$$

## Exercise 2

Show that the geometric multiplicity of an Eigenvalue of the companion matrix of a multistep method, i.e. a matrix like

$$A = \begin{pmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & 0 & 1 \\ -\alpha_0 & -\alpha_1 & \cdots & -\alpha_{m-1} \end{pmatrix}.$$

has to be 1.

**Exercise 3**

Compute the solutions of the explicit and implicit Euler-method of the initial value problem

$$\begin{pmatrix} y_1'(t) \\ y_2'(t) \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \lambda_1 + \lambda_2 & \lambda_1 - \lambda_2 \\ \lambda_1 - \lambda_2 & \lambda_1 + \lambda_2 \end{pmatrix} \begin{pmatrix} y_1(t) \\ y_2(t) \end{pmatrix} \quad \text{with initial value} \quad \begin{pmatrix} y_1(0) \\ y_2(0) \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}.$$

For this you can compute the eigenvalues of the matrix

$$A = \begin{pmatrix} \lambda_1 + \lambda_2 & \lambda_1 - \lambda_2 \\ \lambda_1 - \lambda_2 & \lambda_1 + \lambda_2 \end{pmatrix}$$

and then compute a decomposition of the form  $A = C^{-1}DC$  with a diagonal matrix  $D$ . Derive a formula of the solution of the explicit and the implicit Euler-method. Are the solutions bounded for  $\lambda_1, \lambda_2 < 0$ ?

**Exercise 4**

Verify whether the multistep methods

$$y_{k+2} - 4y_{k+1} + 3y_k = -2\tau f(t_k, y_k)$$

and

$$y_{k+4} - y_k = \frac{h}{3} (8f(t_{k+3}, y_{k+3}) - 4f(t_{k+2}, y_{k+2}) + 8f(t_{k+1}, y_{k+1}))$$

are convergent.